



SUN#DAILY#SPF30 — CPF 4301 Oil-in-Water Sun Cream

## SUN#DAILY#SPF30 - CPF 4301

Oil-in-Water Sun Cream

Phase	Trade name / Supplier	INCI name	Wt %
Α	Water	Water / Aqua	60.75
	SunSpheres™ BIO SPF Booster / Dow	Microcrystalline Cellulose	1.50
	UCON 75-H-450 / Dow	PEG/PPG-17/6 Crosspolymer	0.50
	Butylene Glycol	Butylene Glycol	2.00
	VERSENE™ NA₂ Crystals Chelating Agent / Dow	Disodium EDTA	0.10
	SymSave H / Symrise	Hydroxyacetophenone	0.50
	ACULYN™ 38 Rheology Modifier / Dow	Acrylates / Vinyl Neodecanoate Crosspolymer	3.00
В	Procol CS20D / Making Cosmetics	Cetearyl Alcohol (and) Ceteareth-20	1.75
	Arlacel 165 / Spectrum	Glyceryl Stearate (and) PEG-100 Stearate	2.00
	CCT / Spectrum	Caprylic/Capric Triglyceride	5.00
	C12-15 Alkyl Benzoate / Making Cosmetics	C12-15 Alkyl Benzoate	5.00
	Parsol HMS / DSM	Homosalate	5.00
	Parsol EHS / DSM	Octisalate / Ethylhexyl Salicylate	5.00
	Parsol 1789 / DSM	Avobenzone / Butyl Methoxydibenzoylmethane	3.00
	Parsol 340 / DSM	Octocrylene	4.00
С	TEA, 99% / Dow	Triethanolamine	0.40
D	Neolone PH 100 Preservative / DuPont	Phenoxyethanol	0.50

Disclaimer: Contained in this package is a sample prepared as per the formulation described on this card. Any variation in the formulation/procedure

## Processing instructions:

- Add SunSpheree™ BIO SPF Booster to water and mix until dispersed/no clumps (-30 minutes). Then start heating to 70-75°C and continue mixing. Add Symsaw H, Butylene Glycol, UCON Fluid, VERSENE™ NA, Crystals Chelating Agent, followed by ACULYIN™ 38 Rheology Modifler. Continue mixing and heating to 70-75°C.
- Mix sunscreen ingredients of phase B together and start heating to 70-75°C. Continue mixing/heating until Avobenzone dissolves, then add remainder of phase B ingredients. Mix at 70-75°C until all ingredients are melted/dissolved.
- With phase B @ 70°C add to phase A (70-75°C) and mix well for 5 min.
- Homogenize at 10,000 rpm for 3 min.
  After 2 min, add half of phase C (TEA) and continue to homogenize.
- Switch back to standard mixing and add the other half of phase C. When batch reaches 45°C, add phase D and continue mixing until batch reaches 25°C. Adjust pH to 6,0-7,0 with phase C. if neressary.

Appearance: White cream

Viscosity: 60.000-80.000 cPs

pH: 6.0-7.0

In vivo SPF 37

## Image: AdobeStock 293535811

may cause performance to change.

NOTICE: No freedom from infringement of any patient connect by Dow or offens is to be infermed. Because use conditions and applicable laws may differ from one location to another and may change with firm. Quictorier is responsible for clasterier in such parts and preparation for clasterier is used for formative flat Custorier's violent and disposal practices are in compliance with interpolational to another in the disposal practices are in compliance with interpolational to another interpolation for another

®™ Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

© 2021 The Dow Chemical Company. All rights reserved.